

PRIMAL-DUAL COVARIATE BALANCE AND MINIMAL DOUBLE ROBUSTNESS VIA ENTROPY BALANCING

QINGYUAN ZHAO

Department of Statistics, Stanford University

DANIEL PERCIVAL

Google Inc.

ABSTRACT. Covariate balance is a conventional key diagnostic for methods used estimating causal effects from observational studies. We show covariate balance can also be directly used in the estimation of mean causal effects by studying a recently proposed entropy maximization method called Entropy Balancing (EB). In order to avoid the tautology of repeatedly checking covariate balance, EB exactly matches the covariate moments for the different experimental groups in its optimization problem. We find that the primal and dual problems of EB correspond, respectively, to the implicit linear model of outcome and propensity score logistic regression, with features being the balanced moments. Consequently, we prove EB enjoys some desirable statistical properties: it is doubly robust with respect to the linear models and reaches the asymptotic semiparametric variance bound when both models are correct. Our theoretical results and simulations suggest EB is a very appealing alternative to the conventional propensity score weighting estimators.

1. INTRODUCTION

Consider an observational study, in which one of two conditions (denoted “treatment” and “control”) are not randomly assigned. Such observational data often serve as the basis for many biomedical, economical and social researches, where the researchers wish to study causality between treatment and some response but fail or are not able to run a fully randomized experiment. Making a causal conclusion from observational data is essentially difficult because treatment exposure may be related to some attributes that are also related to the outcome, so the treatment and control groups may be seriously imbalanced in these attributes. If the researcher simply ignores this imbalance, the causal effect estimator could be severely biased.

E-mail addresses: qyzhao@stanford.edu, dancsi@google.com.

Date: August 5, 2015.

Key words and phrases. Causal Inference, Double Robustness, Exponential Tilting, Convex Optimization, Survey Sampling.

We would like to thank Jens Hainmueller, Trevor Hastie, Hera He, William Heavlin, Diane Lambert, Daryl Pregibon, and Jean Steiner for their helpful comments.

To adjust for covariate imbalance, the seminal work of Rosenbaum and Rubin (1983) pointed out the essential role of propensity score, the probability of exposure to treatment conditional on observed covariates. Many statistical methods based on propensity score are subsequently proposed to estimate the mean causal effect. A popular approach is to stratify the observations based on their estimated propensity scores (Rosenbaum and Rubin, 1984). The mean causal effect is then estimated by the weighted average of within-stratum effects. An alternative method (e.g. Hirano and Imbens, 2001) is to construct weights for individual observations by using their estimated propensity scores.

While propensity score stratification is usually more robust to misspecified models, propensity score weighting has more attractive theoretical properties. Hirano et al. (2003) showed that nonparametric propensity score weighting may achieve the semiparametric efficiency bound for the estimation of mean causal effect derived by Hahn (1998). Another desirable property is double robustness. The pioneering work of (Robins et al., 1994) pointed out that propensity score weighting can be further augmented by an outcome regression model. The resulting estimator has the following so-called double robustness property:

Property 1. *If either the propensity score model or the outcome regression model is correctly specified, the mean causal effect estimator is statistically consistent.*

However, propensity score methods commonly suffer from a serious drawback in practice. Often, empirical covariate imbalance exists between the treatment and control groups. Propensity score models often fail to correct for this imbalance. As a result, the estimator of causal effect may be severely biased (Drake, 1993; Smith and Todd, 2001; Kang and Schafer, 2007). A misspecified propensity score model is the most likely root cause, as the true propensity score always will stochastically balance the associated covariates. In order to avoid model misspecification, applied researchers usually increase the complexity of the propensity score model until a sufficiently balanced solution is found. This cyclical process of modeling propensity score and checking covariate balance is sometimes criticized as the “propensity score tautology” (Imai et al., 2008) and, moreover, has no guarantee of even finding a balanced solution.

To avoid repeatedly checking the covariate balance, an alternative is the multivariate matching method based on Mahalanobis distance, which has appealing theoretical properties if covariates have ellipsoidal distributions (Rubin, 1976a,b). However, such assumptions are rarely valid with actual data, so the multivariate matching method may actually make covariate balance worse (Sekhon, 2011). Moreover, matching procedures often discard a large portion of the data and thus lose efficiency.

Recently, some new approaches have been proposed to incorporate covariate balance directly in the procedure, so covariate balance can be satisfied automatically. For instance, Diamond and Sekhon (2013) devised a matching algorithm called Genetic Matching that maximizes the balance of observed covariates between treated and control groups. Imai and Ratkovic (2014) proposed to include covariate balance in the maximum likelihood estimation of propensity score logit model.

In this paper, we study another method for covariate balance called Entropy Balancing (hereafter EB) proposed by Hainmueller (2011). EB is easily interpretable and fast to compute, thus has already gained some popularity in applied fields such as applied econometrics and social sciences (Marcus, 2013; Ferwerda, 2014). In a

nutshell, EB solves an entropy maximization problem under the constraint of exact balance of covariate moments. It puts weights on the control units and can be viewed as an extension to the matching methods which puts equal weights on matched observations.

We show Entropy Balancing is indeed a very attractive alternative to the conventional propensity score weighting methods. We find EB simultaneously fits a logistic regression model for the propensity score and a linear regression model for the outcome. The features of these regression models are the covariate moments being balanced. In Theorem 1, we show EB has a double robustness property: if at least one of the two models are correctly specified, it is consistent for the Population Average Treatment effect for the Treated (PATT), a common quantity of interest in causal inference and survey sampling. Moreover, EB is semiparametric efficient if both models are correctly specified. The two (generalized) linear models have an exact correspondence to the primal and dual optimization problem used to solve EB, revealing an interesting connection between doubly robust estimation and convex optimization.

Since EB does not have any explicit model and the doubly robust property we prove is with respect to the simplest (generalized) linear models, we call EB has a “minimal” double robustness property. Furthermore, the minimality of EB reveals an interesting new relationship between double robustness and covariate balance. Traditionally, one may view the propensity score model and the outcome regression model as contributing independently to the robustness of the causal effect estimator. That is why Robins et al. (1994) named the property “double robustness”. Our study of EB reveals that at least certain forms of the two models are equivalent to each other. The two models essentially produce balanced groups, which reduce the bias of the causal effect estimator. Our understanding of this relationship of covariate balance and double robustness is shown in Figure 1 and is more thoroughly discussed in Section 7.1.




FIGURE 1. The central role of covariate balance in observational study. Dashed arrows: conventional understanding of double robustness. Solid arrows: our understanding of double robustness via entropy balancing.

The rest of the paper is organized as follows. In Section 2, we give a brief review of causal inference and doubly robust estimators. In Section 3 we introduce Entropy Balancing and in Section 4 we discuss the theoretical properties of EB. The empirical performance of EB is studied in Section 5 by simulations. Finally in Section 6 we discuss some extensions of EB and in Section 7 we discuss how propensity scoring and outcome regression are linked by covariate balance.

2. BACKGROUND

First, we review some building blocks of the causal model for non-randomized studies first introduced by Rubin (1974) and the framework of doubly robust estimators. Many of the concepts introduced here are deeply connected to our analysis of EB in the following sections.

2.1. Rubin’s Causal Model. The causal inference problem we consider is best explained by the notion of potential outcome. In Rubin’s causal model, each unit i is associated with a pair of potential outcomes: the response $Y_i(1)$ that is realized if $T_i = 1$ (treated), and another response $Y_i(0)$ realized if $T_i = 0$ (control). Notice that it is impossible to observe both $Y(0)$ and $Y(1)$ on the same unit. This is often referred as the “fundamental problem of causal inference” (Holland, 1986).

We assume the observational units are independent and identically distributed copies from a population, for which we wish to infer the treatment’s effect. In this paper we only consider continuous outcome and focus on the Population Average Treatment effect on the Treated (PATT):

$$(1) \quad \gamma = E[Y(1)|T = 1] - E[Y(0)|T = 1] \triangleq \mu(1|1) - \mu(0|1).$$

This quantity also occurs in survey sampling with missing data (Cassel et al., 1976; Kang and Schafer, 2007). The methods that we consider in this paper also naturally extend to survey sampling problems.

Another common quantity considered in causal inference is the Population Average Treatment Effect (PATE)

$$\tau = E[Y(1)] - E[Y(0)] \triangleq \mu(1) - \mu(0),$$

which is related to PATT through the identity

$$\mu(0) = \mu(0|0)P(T = 0) + \mu(0|1)P(T = 1).$$

The fundamental problem of causal inference is to estimate the “counterfactual” means $\mu(0|1)$ and $\mu(1|0)$. In this paper we focus on the estimation of $\mu(0|1)$.

Along with the treatment exposure T_i and outcome Y_i , each experiment unit i is usually associated with another set of covariates denoted by X_i . In observational studies, both treatment assignment and outcome can be related to the covariates, which may cause serious selection bias. However, as highlighted by the seminal work Rosenbaum and Rubin (1983), it is possible to correct this selection bias under the following two assumptions:

Assumption 1 (strong ignorability). $(Y(0), Y(1)) \perp T \mid X$.

Assumption 2 (overlap). $0 < P(T = 1|X) < 1$.

Intuitively, the first assumption implies that the observed covariates contain all the information that may cause the selection bias, i.e. there is no confounding variable, and the second assumption ensures this bias-correction information is present across the entire domain of X . We will see these two assumptions play a foundational role in the doubly robust estimators (Section 2.2) and the EB method (Section 3 and Section 4.2).

Since the covariates X contain all the confounding information of selection bias, it is important to understand the relationship between T , Y and X . Under Assumption 1 (strong ignorability), the joint distribution of (X, Y, T) is determined by the

marginal distribution of X and two conditional distributions given X . The first conditional distribution $e(X) = P(T = 1|X)$ is often called the propensity score and plays a central role in causal inference. Rosenbaum and Rubin (1983) proved that under Assumptions 1 (strong ignorability) and 2 (overlap) $(Y(0), Y(1)) \perp T | e(X)$, i.e. the propensity score function $e(X)$ itself is sufficient to produce unbiased estimates of causal effect. The second conditional distribution is the density of $Y(0)$ and $Y(1)$ given X . Since we only consider the mean causal effect in this paper, it suffices to study the regression function $E[Y(0)|X]$ and $E[Y(1)|X]$, denoted by $g_0(X)$ and $g_1(X)$.

2.2. Doubly Robust Estimators. Many existing estimators of mean causal effect are special cases of the doubly robust estimation framework (Robins et al., 1994). Readers are recommended to check the subsequent literature (e.g. Robins and Wang, 2000; Bang and Robins, 2005) and the review article by Kang and Schafer (2007) for more details about doubly robust estimators. When using doubly robust estimators, first we need to build individual models for both propensity score $e(X)$ and outcome $(g_0(X), g_1(X))$, then combine the two models with a doubly robust estimator. We review this procedure in this section.

The first component of doubly robust estimators is called inverse-probability weighting (IPW). Suppose we have an estimate of $e(X)$, denoted by $\hat{e}(X)$. For simplicity, we write $\hat{e}(X_i)$ as $\hat{e}_i$. The (weight-normalized) IPW estimators are:

$$(2) \quad \hat{\gamma}^{\text{IPW}} = \sum_{T_i=1} \frac{1}{n_1} Y_i - \sum_{T_i=0} \frac{\hat{e}_i(1 - \hat{e}_i)^{-1}}{\sum_{T_i=0} \hat{e}_i(1 - \hat{e}_i)^{-1}} Y_i.$$

Here $\sum_{T_i=t}$ is defined as summation over all units i such that $T_i = t$. This notation will be repeatedly used throughout the paper.

In this formula, the control units are associated with weights proportional to $\hat{e}_i(1 - \hat{e}_i)^{-1}$ to resemble the full population. The most popular choice of obtaining $\hat{e}(X)$ is via logistic regression, where $\text{logit}(e(X))$ is modeled by $\sum_{j=1}^p \theta_j c_j(X)$ and $c_j(X)$ are functions of the covariates.

The second component of doubly robust estimators is an outcome regression model, which estimates the mean potential outcomes $g_t(X)$ by $\hat{g}_t(X)$, $t = 0, 1$. A common choice of $\hat{g}_t$ is the linear regression model $\hat{g}_t(X) = \sum_{j=1}^p \hat{\beta}_{t,j} c_j(X)$, which leads to the OLS estimators:

$$(3) \quad \hat{\gamma}^{\text{OLS}} = \frac{1}{n_1} \sum_{T_i=1} Y_i - \hat{g}_0(X_i).$$

This approach is also called ‘‘covariance adjustment’’ in the literature (Rosenbaum and Rubin, 1983; Rosenbaum et al., 2002).

The third and last component, a doubly robust semiparametric estimator, combines the first two models using residual bias correction. The following example was first proposed by Cassel et al. (1976) in survey sampling and later generalized by Robins et al. (1994):

$$(4) \quad \hat{\gamma}^{\text{DR}} = \hat{\gamma}^{\text{OLS}} - \sum_{T_i=0} \frac{\hat{e}_i(1 - \hat{e}_i)^{-1}}{\sum_{T_i=0} \hat{e}_i(1 - \hat{e}_i)^{-1}} (Y_i - \hat{g}_0(X_i)).$$

$\hat{\gamma}^{\text{DR}}$ can be viewed as a correction of $\hat{\gamma}^{\text{DR}}$. The bias correction term replaces Y_i in the IPW estimators (2) with the regression residual. The estimator $\hat{\gamma}^{\text{DR}}$ have

the desirable double protection we mentioned earlier and the following efficiency property.

Property 2. *If both propensity score and outcome regression models are correctly specified, the estimator asymptotically achieve the semiparametric variance bound derived in (Hahn, 1998).*

Proposition 1. $\hat{\tau}^{DR}$ and $\hat{\gamma}^{DR}$ satisfy Property 1 and 2.

Proof. Here we only prove the double robustness property and refer the readers to (Rothe and Firpo, 2013) for a proof for efficiency.

In the case that the outcome model is correct specified (i.e. $\hat{g}_0(X) \rightarrow g_0(X)$ as $n \rightarrow \infty$), $\hat{\tau}^{OLS}$ and $\hat{\gamma}^{OLS}$ are already consistent, and the bias correction term in (4) has mean going to 0 as $n \rightarrow \infty$. In the case where the outcome model is misspecified but the propensity scores are correctly specified, the additional term in (4) are consistent estimators for the bias of the OLS estimators. $\square$

Since all models are vulnerable to misspecification, the additional protection DR estimators provide can give the applied researchers much more faith on their causal effect estimates. However, Kang and Schafer (2007) point out the standard DR estimator $\hat{\tau}^{DR}$ could behave poorly when both models are misspecified. Many subsequent doubly robust estimators are proposed to further improve the performance under misspecified models, see Tan (2006); Cao et al. (2009); Tan (2010); Rotnitzky et al. (2012).

The doubly robust property can also be satisfied through weighted regression. We refer the reader to (Kang and Schafer, 2007; Freedman and Berk, 2008) for more details. In Section 4.3 we will construct a doubly robust weighted least squares estimator of γ using Entropy Balancing.

3. ENTROPY BALANCING

In this section, we review the definition of Entropy Balancing (Hainmueller, 2011). We also expose connections of the Entropy Balancing optimization problem with propensity scoring and outcome regression.

The EB procedure generates a set of weights for all the control units that are used to estimate PATT in equation (1). EB operates by maximizing the entropy of the weights under a set of balancing constraints:

$$\begin{aligned}
 (5) \quad & \underset{w}{\text{maximize}} && - \sum_{T_i=0} w_i \log w_i \\
 & \text{subject to} && \sum_{T_i=0} w_i c_j(X_i) = \bar{c}_j(1), \quad j = 1, \dots, p, \\
 & && \sum_{T_i=0} w_i = 1, \\
 & && w_i > 0, \quad i = 1, \dots, n.
 \end{aligned}$$

Hainmueller (2011) refers to the set $\{c_j(\cdot)\}_{j=1}^p$ as moment functions of the covariates. These can be essentially any transformation of X , not necessarily polynomial functions. In Section 4.1 and 4.2, we show in Section 2.2 that they indeed serve the same purposes as linear features in the propensity score model or outcome model. In order to weight the control units to resemble the treatment population, the balancing targets $\bar{c}_j(1)$, $j = 1, \dots, p$ are the empirical “moments” of the treatment

population $\bar{c}_j(1) = \frac{1}{n_1} \sum_{T_i=1} c_j(X_i)$. Throughout the paper we will use $c(X)$ and $\bar{c}(1)$ to stand for the vector of $c_j(X)$ and $\bar{c}_j(1)$, $j = 1, \dots, p$.

Since EB seeks to empirically match the control and treatment covariate distributions, we can draw connections between EB and density estimation. This connection allows us to expose the relationships between EB and propensity score modeling. In particular, the EB optimization problem (5) is a finite sample version of the minimum relative entropy principle in density estimation. Letting $m(x)$ be the density function of the covariates X for the control population, the minimum relative entropy principle estimates the density of the treatment population as

$$(6) \quad \underset{\tilde{m}}{\text{maximize}} \ H(\tilde{m}||m) \quad \text{subject to } E_{\tilde{m}}[c(X)] = \bar{c}(1).$$

Here, $H(\tilde{m}||m) = E_{\tilde{m}}[\log(\tilde{m}(X)/m(X))]$ is the relative entropy between the distribution $\tilde{m}$ and the distribution m . As an estimate of the distribution of the treatment group, the optimal $\tilde{m}$ of (6) is the “closest” to the control distribution among all distributions satisfying the moment constraints.

Let $w(x) = [P(T=1) \cdot \tilde{m}(x)]/[P(T=0) \cdot m(x)]$ be the population version of the inverse probability weights in (2). Applying a change of measure, we can rewrite (6) as an optimization problem over w :

$$(7) \quad \underset{w}{\text{maximize}} \ E_m[w(X) \log w(X)] \quad \text{subject to } E_m[w(X)c(X)] = \bar{c}(1).$$

EB optimization (5) is the a finite sample version of this problem, where the distribution m is replaced by the empirical distribution of the control units.

Next, we show the above heuristic allows us to view EB as a propensity score model. By using the Lagrangian multipliers, we can show the solution to (6) belongs to the family of exponential titled distributions of m (Cover and Thomas, 2012):

$$m_\theta(x) = m(x) \exp(\theta^T c(x) - \psi(\theta)).$$

Here, $\psi(\theta)$ is the moment generating function of this exponential family. Consequently, the solution of (7) is

$$(8) \quad \frac{P(T=1|X=x)}{P(T=0|X=x)} = w(x) = \exp(\alpha + \theta^T c(x))$$

where $\alpha = \log(P(T=1)/P(T=0))$. Equation (8) is exactly the logistic regression model of the propensity score using $c(x)$ as features. This implies EB implicitly fits a logistic regression model.

However, the EB estimate differs from the maximum likelihood estimate of logistic regression. The dual optimization problem of (5) is

$$(9) \quad \underset{\theta}{\text{minimize}} \quad \log \left(\sum_{T_i=0} \exp \left(\sum_{j=1}^p \theta_j c_j(X_i) \right) \right) - \sum_{j=1}^p \theta_j \bar{c}_j(1),$$

which is different from the maximum likelihood problem

$$(10) \quad \underset{\theta}{\text{minimize}} \quad \sum_{i=1}^n \log \left(1 + \exp \left(- (2T_i - 1) \sum_{j=1}^p \theta_j c_j(X_i) \right) \right).$$

The loss functions in (9) and (10) are different. Further, (9) only uses the control units in the minimization.

The unique the solution $\hat{\theta}^{\text{EB}}$ to (9) can be easily computed by Newton method and the EB weights (solution to the primal problem (5)) are immediately given by the Karush-Kuhn-Tucker conditions

$$(11) \quad w_i^{\text{EB}} = \frac{\exp(\sum_{j=1}^p \hat{\theta}_j^{\text{EB}} c_j(X_i))}{\sum_{T_i=0} \exp(\sum_{j=1}^p \hat{\theta}_j^{\text{EB}} c_j(X_i))}, \quad \forall T_i = 0.$$

Hainmueller (2011) proposes to use the weighted average $\sum_{T_i=0} w_i^{\text{EB}} Y_i$ to estimate the counterfactual mean $E[Y(0)|T = 1]$. This gives the Entropy Balancing estimator of PATT

$$(12) \quad \hat{\gamma}^{\text{EB}} = \sum_{T_i=1} \frac{Y_i}{n_1} - \sum_{T_i=0} w_i^{\text{EB}} Y_i.$$

Finally, EB is related to the existing literature that estimates mean causal effect by an empirical likelihood approach (Wang and Rao, 2002; Tan, 2006; Qin and Zhang, 2007; Tan, 2010). EB is different from these methods in two ways. First, no explicit propensity score or outcome regression model is built in the original proposal of EB, while all the other methods rely on at least one of them. Second, instead of the empirical likelihood objective $\sum_{i=1}^n \log w_i$, EB uses Shannon entropy $\sum_{i=1}^n w_i \log w_i$ as the discrepancy function. This makes EB a convex optimization problem, and leads to a logistic propensity score model as we have shown earlier. We can easily generalize EB to other forms of the propensity score model by changing the discrepancy function.

4. PROPERTIES OF ENTROPY BALANCING

Due to its interpretability, Entropy Balancing has already been used in many applications (e.g. Marcus, 2013; Ferwerda, 2014). However, Hainmueller (2011) did not investigate its statistical properties. In this section, we study Entropy Balancing in the framework of Rubin's causal model and doubly robust estimators that are described earlier in Section 2. In particular, we show why Entropy Balancing should be preferred over the conventional maximum likelihood logistic regression for propensity score.

4.1. Existence. We first describe the conditions under which the EB problem (5) admits a solution. The existence of w^{EB} depends on the solvability of the moment matching constraints

$$(13) \quad \sum_{T_i=0} w_i c_j(X_i) = \bar{c}_j(1), \quad j = 1, \dots, p, \quad w > 0, \quad \sum_{T_i=0} w_i = 1.$$

As one may expect, this is closely related to the existence condition of maximum likelihood estimate of logistic regression (Silvapulle, 1981; Albert and Andersen, 1984). An easy way to obtain such condition is through the dual problem of (10)

$$(14) \quad \begin{aligned} & \underset{w}{\text{maximize}} && - \sum_{i=1}^n [w_i \log w_i + (1 - w_i) \log(1 - w_i)] \\ & \text{subject to} && \sum_{T_i=0} w_i c_j(X_i) = \sum_{T_i=1} w_i c_j(X_i), \quad j = 1, \dots, p, \\ & && 0 < w_i < 1, \quad i = 1, \dots, n. \end{aligned}$$

Thus, the existence of $\hat{\theta}^{\text{MLE}}$ is equivalent to the solvability of the constraints in (14), which is the overlap condition first given by Silvapulle (1981).

Intuitively, in the space of $c(X)$, the solvability of (13) or the existence of w^{EB} means there is no hyperplane separating $\{c(X_i)\}_{T_i=0}$ and $\bar{c}(1)$. In contrast, the solvability of (14) or the existence of w^{MLE} means there is no hyperplane separating $\{c(X_i)\}_{T_i=0}$ and $\{c(X_i)\}_{T_i=1}$. Hence the existence of EB requires a stronger condition than the logistic regression MLE.

The next proposition suggests that the existence of w^{EB} and hence w^{MLE} is guaranteed by Assumption 2 (overlap) with high probability.

Proposition 2. *Suppose Assumption 2 (overlap) is satisfied and the expectation of $c(X)$ exist, then $P(w^{\text{EB}} \text{ exists}) \rightarrow 1$ as $n \rightarrow \infty$. Furthermore, $\sum_{i=1}^n (w_i^{\text{EB}})^2 \rightarrow 0$ in probability as $n \rightarrow \infty$.*

The proof of this proposition is deferred to Appendix A.

4.2. Double Linear-Robustness. We now give the main theorem of the paper, which demonstrates that although EB does not contain a propensity score model or a outcome regression model, it nonetheless has a double robustness property.

Theorem 1. *Let Assumption 1 (strong ignorability) and Assumption 2 (overlap) be given. Additionally, assume the expectation of $c(X)$ exists and $\text{Var}(Y(0)) < \infty$. Then Entropy Balancing satisfy Property 1 and 2, in the sense that $\text{logit}(e(X)) = \text{logit}(P(T = 1|X))$ and $g_0(X) = E[Y(0)|X]$ are modeled by linear functions of $c(X)$. The exact statement is*

- (1) *If $\text{logit}(e(X))$ or $g_0(X)$ is linear in $c_j(X)$, $j = 1, \dots, R$, then $\hat{\gamma}^{\text{EB}}$ is statistically consistent.*
- (2) *Moreover, if $\text{logit}(e(X))$, $g_0(X)$ and $g_1(X)$ are all linear in $c_j(X)$, $j = 1, \dots, R$, then $\hat{\gamma}^{\text{EB}}$ reaches the semiparametric variance bound of γ derived in Hahn (1998).*

Notice that this statement is similar to the results in Qin and Zhang (2007). The empirical likelihood (EL) estimator derived therein involves an explicit propensity score model and balancing some auxiliary functions, like the moment functions $c(X)$ in Entropy Balancing. The main conclusion in Qin and Zhang (2007) is that the EL estimator has the same properties as in Theorem 1, but the condition $\text{logit}(e(X))$ is linear in $c(X)$ is replaced by the propensity score model is correctly specified.

The proof of the first claim in Theorem 1 reveals an elegant correspondence between the primal-dual optimization and the statistical property of double robustness, so we sketch the proof here. The proof of the second claim is deferred to Section 4.4.

Proof sketch. The consistency under the linear model of $Y(0)$ is easily obtained using the moment balancing constraints in the primal optimization problem (5). To see this, suppose the true outcome model is

$$(15) \quad g_0(X) = E[Y(0)|X] = \sum_{j=1}^p \beta_j c_j(X)$$

for some β_j and moment functions c_j , $j = 1, \dots, p$. By Assumption 1 (strong ignorability), we have

$$\begin{aligned}
 \mathbb{E}[Y(0)|T=1] &= \mathbb{E}[\mathbb{E}[Y(0)|X]|T=1] \\
 &= \mathbb{E}\left[\sum_{j=1}^p \beta_j c_j(X) \middle| T=1\right] \\
 &= \sum_{j=1}^p \beta_j \mathbb{E}[c_j(X)|T=1].
 \end{aligned}
 \tag{16}$$

Let $\epsilon_i = Y_i(0) - \sum_{j=1}^p \beta_j c_j(X_i)$, so ϵ_i is uncorrelated with $c_j(X_i)$ for $j = 1, \dots, p$. The EB estimator, on the other hand, is

$$\begin{aligned}
 \sum_{T_i=0} w_i Y_i &= \sum_{T_i=0} w_i Y_i(0) \\
 &= \sum_{T_i=0} w_i \left[\sum_{j=1}^p \beta_j c_j(X_i) + \epsilon_i \right] \\
 &= \sum_{j=1}^p \beta_j \sum_{T_i=0} w_i c_j(X_i) + \sum_{T_i=0} w_i \epsilon_i \\
 &= \sum_{j=1}^p \beta_j \bar{c}(1) + \sum_{T_i=0} w_i \epsilon_i.
 \end{aligned}
 \tag{17}$$

Since $\bar{c}(1)$ is unbiased and consistent for $\mathbb{E}[c_j(X)|T=1]$, it is obvious that $\hat{\gamma}^{\text{EB}}$ is always unbiased for γ . Furthermore, because $\sum_i w_i^2 \xrightarrow{P} 0$ by Proposition 2 and $\text{Var}(\epsilon_i) < \infty$, the consistency of EB also follows immediately.

The consistency under the linear model of $\logit(\mathbb{P}(T=1|X))$ is a consequence of the dual optimization problem (9). Intuitively, the dual problem (9) can be viewed as fitting a logistic regression model of the propensity using a loss function different from the usual binomial likelihood, as discussed in 3. In Appendix B we prove this heuristic rigorously using the M-estimation theory. $\square$

Remark 1. The calculations in (16) and (17) implies the difference of the target $\sum_{T_i=0} w_i Y_i$ and the target $\mathbb{E}[Y(0)|T=1]$ is

$$\sum_{T_i=0} w_i Y_i - \mathbb{E}[Y(0)|T=1] = \sum_{j=1}^p \beta_j \left[\sum_{i=1}^n w_i c_j(X_i) - \mathbb{E}[c_j(X)|T=1] \right] + \sum_{T_i=0} w_i \epsilon_i.
 \tag{18}$$

The decomposition (18) only requires the true outcome model is (15). It holds for any weighting estimator, including propensity score matching and stratification which are just special ways of weighting. On the right hand side of (18), the first term is related to the covariate imbalance of the weighting method, while the second term is purely random noise. Thus, for a particular realization of the observational data (X, Y, T) , the role of covariate balance is to minimize the bias of the weighting estimator. Conversely, if the propensity score model does not balance a covariate moment and the moment is correlated with $Y(0)$, that means the estimator is always biased for this realization. When we observe many realization

of the same observational study, this bias will have mean 0 only if the propensity score is correctly specified.

4.3. Entropy Balancing and Outcome Regression. Although the original estimator $\hat{\gamma}^{\text{EB}}$ in (12) does not include an outcome regression model, Theorem 1 shows it still has a double robustness property. In this section we consider the combination of entropy balancing weights and a user-specified outcome regression model.

The heuristic that EB implicitly fits a logistic regression model suggests that it can be used in the doubly robust estimator framework. If we replace the inverse probability weight $\hat{e}_i/(1 - \hat{e}_i)$ in (4) by the EB weight w_i^{EB} in (5), a doubly robust estimator of γ is

$$(19) \quad \hat{\gamma}^{\text{EB-DR}} = \sum_{T_i=1} \frac{1}{n_1} (Y_i - \hat{g}_0(X_i)) - \sum_{T_i=0} w_i^{\text{EB}} (Y_i - \hat{g}_0(X_i)).$$

Corollary 1. $\hat{\gamma}^{\text{EB-DR}}$ is consistent if $\text{logit}(e(X))$ is linear in $\{c_j(X)\}$ or $\hat{g}_0(x)$ is correctly specified for $g_0(x)$.

An important conclusion of our paper is Entropy Balancing also fits a linear outcome model. This is best seen from the following theorem

Theorem 2. If the fitted outcome regression model is $\hat{g}_0(X) = \sum_{j=1}^p \hat{\beta}_j c_j(X)$, whether or not this model is correctly specified, $\hat{\gamma}^{\text{EB-DR}} = \hat{\gamma}^{\text{EB}}$.

Proof. This simple algebraic fact is easily proved by

$$\begin{aligned} \hat{\gamma}^{\text{EB-DR}} - \hat{\gamma}^{\text{EB}} &= \sum_{T_i=0} w_i^{\text{EB}} \sum_{j=1}^p \hat{\beta}_j c_j(X_i) - \frac{1}{n_1} \sum_{T_i=1} \sum_{j=1}^p \hat{\beta}_j c_j(X_i) \\ &= \sum_{j=1}^p \hat{\beta}_j \left(\sum_{T_i=0} w_i^{\text{EB}} c_j(X_i) - \frac{1}{n_1} \sum_{T_i=1} c_j(X_i) \right) \\ &= 0. \end{aligned}$$

□

This theorem gives a new perspective on the moment constraints in (5). Typically, applied researchers are suggested to check the covariate balance in order to make sure the propensity score model is not too biased. However, Theorem 2 indicates that asking exact covariate balance in the finite sample actually fits an implicit outcome regression model. As a result, Entropy Balancing satisfies the doubly robust property stated in Theorem 1.

In principle, we can use this observation to derive a number of doubly robust estimators. More specifically, we can fit any propensity score model with the additional exact moment balancing constraints. The resulting PATT estimator is doubly robust to that propensity score model and the linear outcome regression model. We prefer Entropy Balancing among this type of estimators because it corresponds to the most commonly used logistic regression model.

4.4. Efficiency. In this section, we compute the asymptotic variance of $\hat{\gamma}^{EB}$ using the general M-estimation theory. Let

$$\begin{aligned}\phi_j(X, T; m) &= T(c_j(X) - m_j), \quad j = 1, \dots, p, \\ \psi_j(X, T; \theta, m) &= (1 - T)e^{\sum_{k=1}^p \theta_k c_k(X)}(c_j(X) - m_j), \quad j = 1, \dots, p, \\ \varphi_{1|1}(X, T, Y; \mu(1|1)) &= T(Y - \mu(1|1)), \\ \varphi(X, T, Y; \theta, \mu(1|1), \gamma) &= (1 - T)e^{\sum_{j=1}^p \theta_j c_j(X)}(Y + \gamma - \mu(1|1)),\end{aligned}$$

and $\zeta(X, T, Y; m, \theta, \mu(1|1), \gamma) = (\phi^T, \psi^T, \varphi_{1|1}, \varphi)^T$ be all the estimating equations. The Entropy Balancing estimator $\hat{\gamma}^{EB}$ is the solution to

$$(20) \quad \frac{1}{n} \sum_{i=1}^n \zeta(X_i, T_i, Y_i; m, \theta, \mu(1|1), \gamma) = 0.$$

In Appendix B we use this to prove the consistency of $\hat{\gamma}^{EB}$ when the propensity score is logit linear in $\{c_j(X)\}_{j=1}^p$. Here we compute the asymptotic variance of $\hat{\gamma}^{EB}$ following the approach described in Stefanski and Boos (2002, sec. 2). To state our result, we need to introduce three different kinds of weighted covariance-like functions for arbitrary length p random vectors a_1 and a_2

$$\begin{aligned}H_{a_1, a_2} &= \text{Cov}(a_1, a_2 | T = 1), \\ G_{a_1, a_2} &= E \left[\frac{e(X)}{1 - e(X)} (a_1 - E[a_1 | T = 1])(a_2 - E[a_2 | T = 1])^T | T = 1 \right], \\ K_{a_1, a_2} &= E[(1 - e(X))a_1 a_2^T | T = 1], \\ K_{a_1, a_2}^m &= E[(1 - e(X))a_1 (a_2 - E[a_2 | T = 1])^T | T = 1].\end{aligned}$$

It is obvious that $H \geq K$ and usually $G \geq H$. To make the notation more concise, $c(X)$ will be abbreviated as c and $Y(0)$ as 0 in subscripts. For example, $H_{c,0} = H_{c(X), Y(0)}$, $G_{c,1} = G_{c(X), Y(1)}$ and $K_c = K_{c(X), c(X)}$.

Theorem 3. *Assume the logistic regression model of T is correct, i.e. $\text{logit}(P(T = 1|X))$ is a linear combination of $\{c_j(X)\}_{j=1}^p$. Let $\pi = P(T = 1)$, then we have $\hat{\gamma}^{EB} \xrightarrow{d} N(\gamma, V^{EB}/n)$ and $\hat{\gamma}^{IPW} \xrightarrow{d} N(\gamma, V^{IPW}/n)$ where*

$$(21) \quad V^{EB} = \pi^{-1} \cdot \{H_1 + G_0 - H_{c,0}^T H_c^{-1} (2G_{c,0} - H_{c,0} - G_c H_c^{-1} H_{c,0} + 2H_{c,1})\},$$

$$(22) \quad V^{IPW} = \pi^{-1} \cdot \{H_1 + G_0 - H_{c,0}^T K_c^{-1} (H_{c,0} - 2K_{c,0}^m + 2K_{c,1}^m)\}.$$

The proof of this theorem is in Appendix B. The H , G and K matrices in Theorem 3 can be estimated from the observed data, yielding approximate sampling variances for $\hat{\gamma}^{EB}$ and $\hat{\gamma}^{IPW}$. Alternatively, variance estimates may be obtained via the empirical sandwich method (e.g. Stefanski and Boos, 2002). In practice (particularly in simulations where we compare to a known truth), we find that the empirical sandwich method is more stable than the plug-in method, which is consistent with the suggestion in Lunceford and Davidian (2004) for PATE estimators.

To complete the proof of the second claim in Theorem 1, we compare these variances with the semiparametric variance bound of γ derived by Hahn (1998):

$$(23) \quad V^* = \pi^{-1} \cdot \{H_1 + G_0 - 2H_{E[Y(0)|X], E[Y(1)|X]} - G_{E[Y(0)|X]} + H_{E[Y(0)|X]}\}.$$

Assume $\text{logit}(P(T = 1|X)) = \theta^T c(X)$ and $E[Y(t)|X] = \beta(t)^T c(X)$, $t = 0, 1$. Then

$$\begin{aligned} H_{c,t} &= \text{Cov}(c(X), Y(t)|T = 1) \\ &= \text{Cov}(c(X), \beta(t)^T c(X)|T = 1) \\ &= H_c \beta(t), \text{ for } t = 0, 1. \end{aligned}$$

Similarly, $G_{c,t} = G_c \beta(t)$, $t = 0, 1$. From here it is easy to check V^{EB} and V^* are the same. Since Entropy Balancing reaches the efficiency bound in this case, obviously $V^{\text{EB}} < V^{\text{IPW}}$.

If $\text{logit}(P(T = 1|X)) = \theta^T c(X)$ but $E[Y(t)|X] = \beta(t)^T c(X)$ is not true for some $t = 0, 1$, there is no absolute guarantee that $V^{\text{EB}} < V^{\text{IPW}}$. In practice, the features $c(X)$ in the linear models of Y are almost always correlated with Y . This correlation compensates the slight efficiency loss of not maximizing the likelihood function in logistic regression. As a consequence, the variance V^{EB} in (21) is usually smaller than V^{IPW} in (22). This efficiency advantage of EB over IPW is verified in the next section using simulations.

5. SIMULATIONS

In this section, we use simulated examples to verify the theorems and claims we made earlier. We will also compare EB weighting with IPW (after logistic regression) and the Covariate Balancing Propensity Score (CBPS) proposed by Imai and Ratkovic (2014). See Section 6 for more discussion of CBPS.

5.1. Kang-Schafer Example. First we consider the simulation model in (Kang and Schafer, 2007), which demonstrates that “in at least some settings, two wrong models are not better than one” for the doubly robust estimator (4). The simulated data consists of $\{X_i, Z_i, T_i, Y_i\}, i = 1, \dots, n\}$. X_i and T_i are always observed, Y_i is observed only if $T_i = 1$, and Z_i is never observed. To generate this data set, X_i is distributed as $N(0, I_4)$, Z_i is computed by first applying the following transformation:

$$\begin{aligned} Z_{i1} &= \exp(X_{i1}/2), \\ Z_{i2} &= X_{i2}/(1 + \exp(X_{i1})) + 10, \\ Z_{i3} &= (X_{i1}X_{i3} + 0.6)^3, \\ Z_{i4} &= (X_{i2} + X_{i4} + 20)^2. \end{aligned}$$

Next we normalize each column such that Z_i has mean 0 and standard deviation 1.

In one setting, Y_i is generated by $Y_i = 210 + 27.4X_{i1} + 13.7X_{i2} + 13.7X_{i3} + 13.7X_{i4} + \epsilon_i$, $\epsilon_i \sim N(0, 1)$ and the true propensity scores are $e_i = \text{expit}(-X_{i1} + 0.5X_{i2} - 0.25X_{i3} - 0.1X_{i4})$. In this case, both Y and T can be correctly modeled by (generalized) linear model of the observed covariates X .

In the other settings, at least one of the propensity score model and the outcome regression model is incorrect. In order to achieve this, the data generating process described above is altered such that Y or T (or both) is linear in the unobserved Z instead of the observed X , though the parameters are kept the same.

For each setting (4 in total), we generated 1000 simulated data sets of size $n = 1000$ and apply various methods discussed earlier, including

- (1) IPW, CBPS: the IPW estimator in (2) with propensity score estimated by logistic regression or CBPS;

- (2) EB: the Entropy Balancing estimator in (12);
- (3) IPW+DR, CBPS+DR: the doubly robust estimator in (4) with propensity score estimated by logistic regression or CBPS.

Notice that the correct mean of Y is always 210. The simulation results are shown in Figure 2. The numbers printed at the top of the figure are standard deviations of each method.

The reader may have noticed some unusual facts in this plot. First, the doubly robust estimator “IPW+DR” performs poorly when both models are misspecified (bottom-right panel in Figure 2). In fact, all the three doubly robust methods are worse than just using IPW. Second, the three doubly robust estimators have exactly the same variance if the Y model is correct (top two panels in Figure 2). It seems that how one fits the propensity score model has no impact on the final estimate. This is related to the observation in (Kang and Schafer, 2007) that, in this example, the plain OLS estimate of Y actually outperforms any method involving the propensity score model. Discussion articles such as Robins et al. (2007) and Ridgeway and McCaffrey (2007) find this phenomenon very uncommon in practice and is most likely due to the estimated inverse probability weights are highly variable, which is a bad setting for doubly robust estimators.

Although the Kang-Schafer example is artificial and is arguably not very likely to occur in practice, we make two comments here about entropy balancing in this unfavorable setting for doubly robust estimators:

- (1) If both T and Y models are misspecified, EB has smaller bias than the conventional “IPW+DR” or “CBPS+DR”. So EB seems to be less affected by such unfavorable setting.
- (2) When T model is correct but Y model is wrong (bottom-left panel in Figure 2), EB has the smallest variance among all estimators. This supports the conclusion of our efficiency comparison of IPW and EB in Section 4.4.

Finally we want to notice that the same simulation setting is used in Tan (2010) to study the performance of a number of doubly robust estimators. The reader can compare the Figure 2 with the results there. Overall, the performance of Entropy Balancing is comparable to the best estimator in Tan (2010).

5.2. Lunceford-Davidian Example. Here we consider another comprehensive simulation example, which is used by Lunceford and Davidian (2004) to study estimators of PATE τ . We find this example is suitable to investigate estimators of PATT γ as well.

In this simulation, the data still consists of $\{\{X_i, Z_i, T_i, Y_i\}, i = 1, \dots, n\}$, but all of them are observed. Both X_i and Z_i are three dimensional vectors. The propensity score is only related to X through:

$$\text{logit}(P(T_i = 1)) = \beta_0 + \sum_{j=1}^3 \beta_j X_{ij}.$$

Note the above does not involve elements of Z_i . The response Y is generated according to

$$Y_i = \nu_0 + \sum_{j=1}^3 \nu_j X_{ij} + \nu_4 T_i + \sum_{j=1}^3 \xi_j Z_{ij} + \epsilon_i; \quad \epsilon_i \sim N(0, 1).$$




FIGURE 2. Results of the Kang-Schafer Example. The methods are: Inverse Propensity Weighting (IPW), Covariate Balancing Propensity Score (CBPS), Entropy Balancing (EB), and doubly robust versions of the first two (IPW+DR, CBPS+DR). Both propensity score model and outcome regression model can be correct or incorrect, so there are four scenarios in total. We generate 1000 simulations of 1000 sample from the example in (Kang and Schafer, 2007) and apply five different estimators. Target mean is 210 and is marked as a black horizontal line to compare the biases of the methods. Numbers printed at $Y = 230$ are the sample standard deviation of each method, in order to compare their efficiency.

The parameters here are set to be

$$\nu = (0, -1, 1, -1, 2)^T;$$

β is set as:

$$\begin{aligned}\beta^{\text{no}} &= (0, 0, 0, 0)^T, \\ \beta^{\text{moderate}} &= (0, 0.3, -0.3, 0.3)^T, \text{ or} \\ \beta^{\text{strong}} &= (0, 0.6, -0.6, 0.6)^T.\end{aligned}$$

The choice of β depends on the level of association of T and X . ξ is based on a similar choice on the level of association of Y and Z :

$$\begin{aligned}\xi^{\text{no}} &= (0, 0, 0)^T, \\ \xi^{\text{moderate}} &= (-0.5, 0.5, 0.5)^T, \text{ or} \\ \xi^{\text{strong}} &= (-1, 1, 1)^T.\end{aligned}$$

The joint distribution of (X_i, Z_i) is specified by taking $X_{i3} \sim \text{Bernoulli}(0.2)$ and then generate Z_{i3} as Bernoulli with

$$P(Z_{i3} = 1|X_{i3}) = 0.75X_{i3} + 0.25(1 - X_{i3}).$$

Conditional on X_{i3} , $(X_{i1}, Z_{i1}, X_{i2}, Z_{i2})$ is then generated as multivariate normal $N(a_{X_{i3}}, B_{X_{i3}})$, where $a_1 = (1, 1, -1, -1)^T$, $a_0 = (-1, -1, 1, 1)^T$ and

$$B_0 = B_1 = \begin{pmatrix} 1 & 0.5 & -0.5 & -0.5 \\ 0.5 & 1 & -0.5 & -0.5 \\ -0.5 & -0.5 & 1 & 0.5 \\ -0.5 & -0.5 & 0.5 & 1 \end{pmatrix}.$$

The data generating process implies the true PATT is $\gamma = 2$. Since the outcome Y depends on both X and Z , we always fit a full linear model of Y using X and Z , if such model is needed. T only depends on X , so it is not necessary to include Z in propensity score modeling. However, as pointed out by Lunceford and Davidian (2004, Sec. 3.3), it is actually beneficial to “overmodel” the propensity score by including Z in the model. Here we will try both possibilities, the “full” modeling of T using both X and Z , and the “partial” modeling of T using only X .

We generated 1000 simulated data sets and the results are shown in Figure 3 for “full” propensity score modeling and Figure 4 for “partial” propensity score modeling. We make the following comments about these two plots:

- (1) IPW and all the other estimators are always consistent, no matter what level of association is specified. This is because the propensity score model is always correctly specified.
- (2) When using full propensity score modeling, all the doubly robust estimators (EB, IPW+DR, CBPS+DR and EB+DR) have almost the same sample variance. This is because all of them are asymptotically efficient.
- (3) CBPS, to our surprise, does not perform very well in this simulation. It has smaller variance than IPW but this comes with the price of some bias. If we use the partial propensity score model (only involve X , Figure 4), this bias is a little smaller but still not negligible. While it is not clear what causes this bias, one possible reason is that the optimization problem of CBPS is nonconvex, so the local solution which is used to construct γ estimator could be far from the global solution. Another possibility is that CBPS uses GMM or Empirical Likelihood to combine likelihood with imbalance penalty, which is less efficient than maximum likelihood directly. Thus, although the estimator is asymptotically unbiased, the convergence spend




FIGURE 3. Results of the Lunceford-Davidian example (full propensity score modeling). The propensity score model and outcome regression model, if applies, are always correctly specified, but the level of association between T or Y with X or Z could be different, ended up with 9 different scenarios. X are confounding covariates and Z only affects the outcome. We generate 1000 simulations of 1000 in each scenario and apply five different estimators. Target PATE is 2 and is marked as a black horizontal line to compare the biases of the methods. Numbers printed at $Y = 5$ are the sample standard deviation of each method, in order to compare their efficiency.

to the true γ is quite slower than IPW. CBPS combined with outcome regression (CB+DR) fixes the bias and inefficiency issue occurred in CBPS without outcome regression.




FIGURE 4. Results of the Lunceford-Davidian example (partial propensity score modeling). The settings are exactly the same as Figure 3 except the methods here don't use Z in their propensity score models.

- (4) EB, in contrast, performs quite well in this simulation. It has relatively small variance overall, especially if we use the “full” model, i.e. both X and Z are being balanced in (5).
- (5) The difference between EB and EB+DR is that while EB only balances “partial” or “full” covariates, EB+DR additionally combines a outcome linear regression model on all the covariates. As proved in Theorem 2, when the “full” covariates are used, EB is exactly the same as EB+DR. We can observe this from Figure 3. When EB only balances “partial” covariates, the two methods are different and indeed EB+DR is more efficient in Figure 4 since it fits the correct Y model.
- (6) Using the “full” propensity score model improves the efficiency of pure weighting estimators (IPW, CBPS and EB) a lot, but has very little impact

on estimators that involves an outcome regression model (IPW+DR and CBPS+DR) compared to "partial" propensity score modeling. Although EB could be viewed as fitting a outcome model implicitly, the "partial" EB estimator only uses X in the outcome model, that is precisely the reason why it is not efficient. Thus there are both robustness and efficiency reasons that one should include all relevant covariates in EB, even if the covariates affect only one of T and Y .

6. EXTENSIONS

6.1. Weighted regression. The weights generated by EB can be used in a weighted outcome regression model

$$(24) \quad \hat{\beta}^{\text{EB-WLS}} = \left(\sum_{T_i=0} w_i^{\text{EB}} X_i X_i^T \right)^{-1} \left(\sum_{T_i=0} w_i^{\text{EB}} X_i Y_i \right),$$

$$(25) \quad \hat{\gamma}^{\text{EB-WLS}} = \frac{1}{n_1} \sum_{T_i=1} (Y_i - X_i \hat{\beta}^{\text{EB-WLS}}).$$

The weighted regression estimator $\hat{\gamma}^{\text{EB-WLS}}$ has the same double robustness property as $\hat{\gamma}^{\text{EB}}$ because it fits a linear model on Y . In general, unlike $\hat{\gamma}^{\text{EB-DR}}$ as in Theorem 2, $\hat{\gamma}^{\text{EB-WLS}}$ is not the same as the original estimator $\hat{\gamma}^{\text{EB}}$.

6.2. Relaxed balancing. We next propose a relaxation of Entropy Balancing and compare it to a recently proposed robust method of fitting propensity scores (Imai and Ratkovic, 2014).

One concern about Entropy Balancing is that the exact moment balancing constraints could be too stringent, especially if we have a lot of them. This could happen in high dimensional problems or if we wish to balance many interactions. In such a case, the solution to the EB problem (5) might not exist. Even if does, the weights could have very high variance. Thus, we propose an extension to optimization problem in (5) with relaxed constraints:

$$(26) \quad \begin{aligned} & \text{minimize}_w \quad \sum_{T_i=0} w_i \log w_i + \frac{1}{2\lambda} \sum_{j=1}^p \left[\sum_{T_i=0} w_i c_j(X_i) - \bar{c}(1) \right]^2 \\ & \text{subject to} \quad \sum_{T_i=0} w_i = 1, \\ & \quad \quad \quad w_i \geq 0, \quad i = 1, \dots, n. \end{aligned}$$

It is obvious that if $\lambda = 0$, (26) is equivalent to (5). The dual problem of (26) is

$$(27) \quad \text{minimize}_\theta \quad \log \left\{ \sum_{T_i=0} \exp \left(\sum_{j=1}^p \theta_j (c_j(X_i) - \bar{c}_j(1)) \right) \right\} + \frac{\lambda}{2} \|\theta\|_2^2.$$

This means the "softened" EB essentially fits a logistic model by minimizing l_2 -regularized exponential loss on half of the data. Regularization is known to improve estimation and prediction accuracy by reducing variance. However, this reduction comes with a price of adding bias to the statistical inference. Since $\lambda = 0$ in

(26) corresponds to (5), EB employs no regularization (27). Therefore, it gives an unbiased estimator of mean causal effect, as we have already seen from Theorem 1.

Recently, Imai and Ratkovic (2014) proposed Covariate Balancing Propensity Score (CBPS), which is related to this relaxation of EB. CBPS explicitly builds a logistic propensity score model and tries to make it robust by combining covariate balancing constraints. However, this robust approach is different from the dual optimization (27) in the following ways:

- (1) CBPS replaces the “half” exponential loss in (27) with the full logistic likelihood loss.
- (2) Instead of the norm of θ , CBPS penalizes by the covariate imbalance induced by the fitted propensity score model, similar to the soft balancing term in the EB primal problem (26).
- (3) CBPS combines the likelihood loss in point (1) and the covariate imbalance in point (2) using Generalized Method of Moments or Empirical Likelihood, two common approaches for solving general estimating equations.

Our simulations in 5 show that EB usually performs better than CBPS. We make two additional notes about CBPS here: first, since it is purely a robust way to fit the propensity score model and does not involve any modeling of Y , it can be combined with an outcome regression model to achieve double robustness (see Section 2.2); second, unlike Entropy Balancing, the optimization problem CBPS proposes is non-convex, has no guarantee of finding the global optimum, and is much slower to solve in practice.

7. CONCLUDING REMARKS

7.1. The role of covariate balance. As a concluding remark, we argue that the two arms of double robustness are not completely independent. Instead, they are collaborating or even competing towards the same goal—covariate balance. This connection gives a new view on using covariate balance as a diagnostic for propensity scoring.

First, suppose we are given a propensity score model that already balances some covariate moments. Theorem 2 indicates that “double-robustifying” the corresponding IPW estimator by an outcome linear regression model on those moments does not change the estimator at all. In other words, this outcome regression model is automatically built-in when we ask for covariate balance in propensity scores.

Second, a linear regression model of the outcome also balances the covariates. Consider a linear regression of the potential outcome $Y(0)$ on the covariate moments $c_j(X)$, $j = 1, 2, \dots, p$. Let X^t denote the matrix $X_{ij}^t = c_j(X_i)$ for i in the group $t = 0$ or 1 and Y^t denote the vector of outcomes in group $t = 0$ or 1 . An OLS estimate of the coefficients is

$$\hat{\beta} = [(X^0)^T X^0]^{-1} (X^0)^T Y^0.$$

The OLS estimator (3) of $E[Y(0)|T = 1]$ is simply

$$\frac{1}{n_1} 1^T (X^1 \hat{\beta}) = \frac{1}{n_1} 1^T \{X^1 [(X^0)^T X^0]^{-1} (X^0)^T\} Y^0,$$

which can be thought as a weighted sum of Y^0 . These weights, when applied to the covariates of group 0, automatically match the unweighted average of group 1.

That is, if we let w_i be the i -th element of $\frac{1}{n_1}1^T \{X^1[(X^0)^T X^0]^{-1}(X^0)^T\}$ for i in group 0, then

$$\sum_{T_i=0} w_i c_j(X_i) = \frac{1}{n_1} \sum_{T_i=1} c_j(X_i),$$

which is exactly the same as the moment balancing constraints in (5). The derivation here implies, surprisingly, that the balancing weights can be also obtained by outcome regression.

For the two reasons above, both propensity scoring and outcome regression are trying improve the covariate balance as illustrated in Figure 1. Further, a logistic regression model of propensity scores and a linear regression model of outcome exhibit a certain degree of equivalence.

7.2. Conclusions. Entropy Balancing is motivated by and directly addresses the “propensity score tautology” of repeatedly checking covariate balance. When a researcher decides to check the covariate balance after modeling propensity scores, he or she is implicitly checking for robustness with respect to certain outcome models (the linear models on those covariate moment she checks). The researcher is also forfeiting robustness to all other outcome models (see Remark 1), unless he or she is truly confident about her propensity score model (for example, only 5 degrees of freedom are used in the propensity score model and 100 target covariate moments are balanced). By using the principle of relative entropy minimization, Entropy Balancing is the “minimal” model for this purpose and should be preferred for its simplicity as well as the strong statistical properties shown in this paper.

APPENDIX A. PROOF OF PROPOSITION 2

Since the expectation of $c(X)$ exist, the weak law of large number says $\bar{c}(1) \xrightarrow{P} \bar{c}^*(1) = E[c(X)|T=1]$. Therefore

Lemma 1. *For any $\epsilon > 0$, $P(\|\bar{c}(1) - \bar{c}^*(1)\|_\infty \geq \epsilon) \rightarrow 0$ as $n \rightarrow \infty$.*

Now condition on $\|\bar{c}(1) - \bar{c}^*(1)\|_\infty \geq \epsilon$, i.e. $\bar{c}(1)$ is in the box of side length 2ϵ centered at $\bar{c}^*(1)$, we want to prove that with probability going to 1 there exists w such that $w_i > 0$, $\sum_{T_i=0} w_i = 1$ and $\sum_{T_i=0} w_i c(X_i) = \bar{c}(1)$. Equivalently, this is saying the convex hull generated by $\{c(X_i)\}_{T_i=0}$ contains $\bar{c}(1)$. We will indeed prove a little stronger result:

Lemma 2. *With probability going to 1 the convex hull generated by $\{c(X_i)\}_{T_i=0}$ contains the box $B_\epsilon(\bar{c}^*(1)) = \{c(x) : \|c(x) - \bar{c}^*(1)\|_\infty \leq \epsilon\}$ for some $\epsilon > 0$.*

Proposition 2 follow immediately from Lemma 1 and Lemma 2. Now we prove Lemma 2. Denote the sample space of X by $\Omega(X)$. Assumption 2 (overlap) implies $\bar{c}^*(1)$ hence $B_\epsilon(\bar{c}^*(1))$ is in the interior of the convex hull of $\Omega(X)$ for sufficiently small ϵ . Let R_i , $i = 1, \dots, 3^p$, be the 3^p boxes centered at $\bar{c}^*(1) + \frac{3}{2}\epsilon b$, where $b \in \mathbb{R}^p$ is a vector that each entry can be $-1, 0$, or 1 . It is easy to check that the sets R_i are disjoint and the convex hull of $\{x_i\}_{i=1}^{3^p}$ contains $B_\epsilon(\bar{c}^*(1))$ if $x_i \in R_i$, $i = 1, \dots, 3^p$.

Since $0 < P(T = 0|X) < 1$, $\rho = \min_i P(X \in R_i|T = 0) > 0$. This implies

$$\begin{aligned}
 (28) \quad P(\exists X_i \in R_i \text{ and } T_i = 0, \forall i = 1, \dots, 3^p) &\geq 1 - \sum_{i=1}^{3^p} P(X \notin R_i|T = 0)^n \\
 &\geq 1 - 3^p(1 - \rho)^n \\
 &\rightarrow 1
 \end{aligned}$$

as $n \rightarrow \infty$. This proves the lemma because the event in the left hand side implies the convex hull generated by $\{c(X_i)\}_{T_i=0}$ contains the desired box. Note that (28) also tells us how many samples we actually need to ensure the existence of w^{EB} . Indeed if $n \geq \rho^{-1}(p \log 2 + \log \delta^{-1}) \geq \log_{(1-\rho)}(\delta 2^{-p})$, then the probability in (28) is greater than $1 - \delta$. Usually we expect $\delta = O(3^{-p})$. If this is the case, the number of samples needed is $n = O(p \cdot 3^p)^1$.

Now we turn to the second claim of the proposition, i.e. $\sum_{T_i=0} w_i^2 \xrightarrow{P} 0$. To prove this, we only need to find a sequence (with respect to growing n) of feasible solutions to (5) such that $\max_i w_i \rightarrow 0$. This is not hard to show, because the probability in (28) is exponentially decaying as n increases. We can pick $n_1 \geq N(\delta, p, \rho)$ such that the probability of the convex hull of $\{x_i\}_{i=1}^{n_1}$ contains $B_\epsilon(\bar{c}^*(1))$ is at least $1 - \delta$, then pick $n_{i+1} \geq n_i + 3^i N(\delta, p, \rho)$ so the convex hull of $\{x_i\}_{i=n_i+1}^{n_{i+1}}$ contains $B_\epsilon(\bar{c}^*(1))$ with probability at least $1 - 3^i \delta$. This means for each $\{x_i\}_{i=n_i+1}^{n_{i+1}}$, $i = 0, 1, \dots$, we have a set of weights $\{\tilde{w}_i\}_{i=n_i+1}^{n_{i+1}}$ such that $\sum_{i=n_i+1}^{n_{i+1}} \tilde{w}_i x_i = \bar{c}(1)$. Now suppose $n_k \leq n < n_{k+1}$, the choice $w_i = \tilde{w}_i/k$ if $i \leq n_k$ and $w_i = 0$ if $i > n_k$ satisfies the constraints and $\max_i w_i \leq k$. As $n \rightarrow \infty$, this implies $\max_i w_i \rightarrow 0$ and hence $\sum_i w_i^2 \rightarrow 0$ with probability tending to 1.

APPENDIX B. PROOF OF THEOREM 1 AND 3

The first claim of Theorem 1 is immediately proved by the following lemma:

Lemma 3. *Under the assumptions in Theorem 1 and suppose $\logit(P[T = 1|X]) = \sum_{j=1}^p \theta_j^* c_j(X)$, then as $n \rightarrow \infty$, $\hat{\theta}^{\text{EB}} \xrightarrow{P} \theta^*$. As a consequence,*

$$E \left[\sum_{T_i=0} w_i^{\text{EB}} Y_i \right] \xrightarrow{P} E[Y(0)|T = 1].$$

Proof. The proof is a standard application of M-estimation (more precisely Z-estimation) theory. We will follow the estimating equations approach described in (Stefanski and Boos, 2002) to derive consistency of $\hat{\theta}^{\text{EB}}$. First we note the first order optimality condition of (9) is

$$(29) \quad \sum_{i=1}^n (1 - T_i) e^{\sum_{k=1}^p \theta_k c_k(X_i)} (c_j(X_i) - \bar{c}_j(1)) = 0, \quad j = 1, \dots, R.$$

We can rewrite (29) as estimating equations. Let $\phi_j(X, T; m) = T(c_j(X) - m_j)$, $j = 1, \dots, R$ and $\psi_j(X, T; \theta, m) = (1 - T) \exp\{\sum_{k=1}^p \theta_k c_k(X)\} (c_j(X) - m_j)$, then (29)

¹Note that this naive rate can actually be greatly improved by Wendel's theorem in geometric probability theory.

is equivalent to

$$(30) \quad \begin{aligned} \sum_{i=1}^n \phi_j(X_i, T_i; m) &= 0, \quad j = 1, \dots, R, \\ \sum_{i=1}^n \psi_j(X_i, T_i; \theta, m) &= 0, \quad j = 1, \dots, R. \end{aligned}$$

Since $\phi(\cdot)$ and $\psi(\cdot)$ are all smooth functions of θ and m , all we need to verify is that $m_j^* = E[c_j(X)|T=1]$ and θ^* is the unique solution to the population version of (30). It is obvious that m^* is the solution to $E[\phi_j(X, T; m)] = 0$, $j = 1, \dots, R$. Now take conditional expectation of ψ_j given X :

$$\begin{aligned} E[\psi_j(X, T; \theta, m^*) | X] &= (1 - e(X))e^{\sum_{k=1}^p \theta_k c_k(X)} (c_j(X) - m_j^*) \\ &= \left(1 - \frac{e^{\sum_{k=1}^p \theta_k^* c_k(X)}}{1 + e^{\sum_{k=1}^p \theta_k^* c_k(X)}}\right) e^{\sum_{k=1}^p \theta_k c_k(X)} (c_j(X) - m_j^*) \\ &= \frac{e^{\sum_{k=1}^p \theta_k c_k(X)}}{1 + e^{\sum_{k=1}^p \theta_k^* c_k(X)}} (c_j(X) - E[c_j(X)|T=1]). \end{aligned}$$

The only way to make $E[\psi_j(X, T; \hat{\theta}, m^*)] = 0$ is to have

$$\frac{e^{\sum_{k=1}^p \hat{\theta}_k c_k(X)}}{1 + e^{\sum_{k=1}^p \theta_k^* c_k(X)}} = \text{const} \cdot P(T=1|X),$$

i.e. $\hat{\theta} = \theta^*$. This proves the consistency of $\hat{\theta}^{\text{EB}}$.

The consistency of $\hat{\gamma}^{\text{EB}}$ is proved by noticing

$$w_i^{\text{EB}} = \frac{\exp(\sum_{j=1}^p \hat{\theta}_j^{\text{EB}} c_j(X_i))}{\sum_{T_i=0} \exp(\sum_{j=1}^p \hat{\theta}_j^{\text{EB}} c_j(X_i))} \xrightarrow{P} \frac{P(T_i=1|X_i)}{1 - P(T_i=1|X_i)},$$

which is the IPW-NR weight defined in (2). $\square$

The second claim is a corollary of Theorem 3, which is proved below. For simplicity we denote $\xi = (m^T, \theta^T, \mu(1|1), \gamma)^T$ and the true parameter as ξ^* . Throughout this section we assume $\text{logit}(e(X)) = \sum_{j=1}^p \theta_j^* c_j(X)$. Denote $\tilde{c}(X) = c(X) - \bar{c}^*(1)$, $e^*(X) = e(X; \theta^*)$, $l^*(X) = \exp\{\sum_{j=1}^p \theta_j^* c_j(X)\} = e^*(X)/(1 - e^*(X))$.

There are two forms of “information” matrix that need to be computed. The first is

$$\begin{aligned} A^{\text{EB}}(\xi^*) &= E \left[-\frac{\partial}{\partial \xi^T} \zeta(X, T, Y; \xi^*) \right] \\ &= \left(E \left[-\frac{\partial}{\partial m^T} \zeta(\xi^*) \right] \quad E \left[-\frac{\partial}{\partial \theta^T} \zeta(\xi^*) \right] \quad E \left[-\frac{\partial}{\partial \mu(1|1)} \zeta(\xi^*) \right] \quad E \left[-\frac{\partial}{\partial \gamma} \zeta(\xi^*) \right] \right) \\ &= E \left[\begin{pmatrix} T \cdot I_R & 0 & 0 & 0 \\ (1-T)l^*(X) \cdot I_R & -(1-T)l^*(X)(c(X) - \bar{c}^*(1))c(X)^T & 0 & 0 \\ 0^T & 0^T & T & 0 \\ 0 & -(1-T)l^*(X)(Y(0) - \mu^*(0|1))c(X)^T & (1-T)l^*(X) & -(1-T)l^*(X) \end{pmatrix} \right] \\ &= \pi \cdot \begin{pmatrix} I_R & 0 & 0 & 0 \\ I_R & -\text{Cov}[c(X)|T=1] & 0 & 0 \\ 0^T & 0^T & 1 & 0 \\ 0 & -\text{Cov}(Y(0), c(X)|T=1) & 1 & -1 \end{pmatrix}. \end{aligned}$$

A very useful identity in the computation of the expectation is

$$\begin{aligned} \mathbb{E}[f(X, Y)|T = 1] &= \pi^{-1} \mathbb{E}[e(X)f(X, Y)] \\ &= \frac{\mathbb{P}(T = 0)}{\pi} \cdot \mathbb{E} \left[\frac{e(x)}{1 - e(X)} f(X, Y) \middle| T = 0 \right]. \end{aligned}$$

The second information matrix is the covariance of $\zeta(X, T, Y; \xi^*)$. Denote $\tilde{Y}(t) = Y(t) - \mu^*(t|1)$, $t = 0, 1$

$$\begin{aligned} B^{\text{EB}}(\xi^*) &= \mathbb{E}[\zeta(X, Y, T; \xi^*)\zeta(X, Y, T; \xi^*)^T] \\ &= \mathbb{E} \left[\begin{pmatrix} T\tilde{c}(X)\tilde{c}(X)^T & 0 & T\tilde{Y}(1)\tilde{c}(X) & 0 \\ 0 & (1-T)l^*(X)^2\tilde{c}(X)\tilde{c}(X)^T & 0 & (1-T)l^*(X)^2\tilde{Y}(0)\tilde{c}(X) \\ T\tilde{Y}(1)\tilde{c}(X)^T & 0^T & T\tilde{Y}^2(1) & 0 \\ 0 & (1-T)l^*(X)^2\tilde{Y}(0)\tilde{c}(X)^T & 0 & (1-T)l^*(X)^2\tilde{Y}^2(0) \end{pmatrix} \right]. \end{aligned}$$

The asymptotic distribution of $\hat{\gamma}^{\text{EB}}$ is $N(\gamma, V^{\text{EB}}(\xi^*)/n)$ where $V^{\text{EB}}(\xi^*)$ is the bottom right entry of $A^{\text{EB}}(\xi^*)^{-1}B^{\text{EB}}(\xi^*)A^{\text{EB}}(\xi^*)^{-T}$. Let's denote

$$H_{a_1, a_2} = \text{Cov}(a_1, a_2|T = 1),$$

$$G_{a_1, a_2} = \mathbb{E} [l^*(X)(a_1 - \mathbb{E}[a_1|T = 1])(a_2 - \mathbb{E}[a_2|T = 1])^T | T = 1]^T,$$

and $H_a = H_{a,a}$, $G_a = G_{a,a}$. So

$$A^{\text{EB}}(\xi^*) = \pi \cdot \begin{pmatrix} I_R & 0 & 0 & 0 \\ I_R & -H_{c(X)} & 0 & 0 \\ 0^T & 0^T & 1 & 0 \\ 0 & -H_{Y(0), c(X)} & 1 & -1 \end{pmatrix},$$

$$A^{\text{EB}}(\xi^*)^{-1} = \pi^{-1} \cdot \begin{pmatrix} I_R & 0 & 0 & 0 \\ H_{c(X)}^{-1} & -H_{c(X)}^{-1} & 0 & 0 \\ 0^T & 0^T & 1 & 0 \\ -H_{c(X), Y(0)}^T H_{c(X)}^{-1} & H_{c(X), Y(0)}^T H_{c(X)}^{-1} & 1 & -1 \end{pmatrix},$$

and

$$B^{\text{EB}}(\xi^*) = \pi \cdot \begin{pmatrix} H_{c(X)} & 0 & H_{c(X), Y(1)} & 0 \\ 0 & G_{c(X)} & 0 & G_{c(X), Y(0)} \\ H_{Y(1), c(X)} & 0^T & H_{Y(1)} & 0 \\ 0 & G_{Y(0), c(X)}^T & 0 & G_{Y(0)} \end{pmatrix}.$$

Thus

$$V^{\text{EB}} = \pi^{-1} \cdot \{H_{c,0}^T H_c^{-1} (H_{c,0} + G_c H_c^{-1} H_{c,0} - 2G_{c,0} - 2H_{c,1}) + H_1 + G_0\}.$$

It would be interesting to compare $V^{\text{EB}}(\xi^*)$ with $V^{\text{IPW}}(\xi^*)$, the asymptotic variance of $\hat{\gamma}^{\text{IPW}}$. The IPW PATT estimator (2) is equivalent to solving the following estimating equations

$$\begin{aligned} \sum_{i=1}^n \left(T_i - \frac{1}{1 + e^{-\sum_{k=1}^p \theta_k c_k(X_i)}} \right) c_j(X_i) &= 0, \quad r = 1, \dots, R, \\ \frac{1}{n} \sum_{i=1}^n \varphi_{1|1}(X_i, T_i, Y_i; \theta, \mu(1|1), \gamma) &= 0, \\ \frac{1}{n} \sum_{i=1}^n \varphi(X_i, T_i, Y_i; \theta, \mu(1|1), \gamma) &= 0. \end{aligned}$$

If we call $K_{a_1, a_2} = E[(1 - e(X))a_1 a_2^T | T = 1]$, we have

$$\begin{aligned} A^{\text{IPW}}(\xi^*) &= E \left[\begin{pmatrix} e^*(X)(1-e^*(X))c(X)c(X)^T & 0 & 0 \\ 0^T & T & 0 \\ -(1-T)l^*(X)\tilde{Y}(0)c(X)^T & (1-T)l^*(X) & -(1-T)l^*(X) \end{pmatrix} \right] \\ &= \pi \cdot \begin{pmatrix} K_{c(X)} & 0 & 0 \\ 0^T & 1 & 0 \\ -H_{Y(0), c(X)} & 1 & -1 \end{pmatrix}. \\ A^{\text{IPW}}(\xi^*)^{-1} &= \pi^{-1} \cdot \begin{pmatrix} K_{c(X)}^{-1} & 0 & 0 \\ 0^T & 1 & 0 \\ -H_{Y(0), c(X)} K_{c(X)}^{-1} & 1 & -1 \end{pmatrix}. \end{aligned}$$

Let $q^*(X) = e^*(X)l^*(X)$,

$$\begin{aligned} B^{\text{IPW}}(\xi^*) &= E \left[\begin{pmatrix} (T-e^*(X))^2 c(X)c(X)^T & T(T-e^*(X))\tilde{Y}(1)c(X) & -(1-T)q^*(X)\tilde{Y}(0)c(X) \\ T(T-e^*(X))\tilde{Y}(1)c(X)^T & T\tilde{Y}^2(1) & 0 \\ -(1-T)q^*(X)\tilde{Y}(0)c(X) & 0 & (1-T)l^*(X)^2 \tilde{Y}^2(0) \end{pmatrix} \right] \\ &= \pi \cdot \begin{pmatrix} K_{c(X)} & K_{c(X), \tilde{Y}(1)} & K_{c(X), \tilde{Y}(0)} - H_{c(X), Y(0)} \\ K_{c(X), \tilde{Y}(1)}^T & H_{Y(1)} & 0 \\ K_{c(X), \tilde{Y}(0)}^T - H_{c(X), Y(0)}^T & 0 & G_{Y(0)} \end{pmatrix}. \end{aligned}$$

V^{IPW} can thus be computed consequently and we omit the details.

REFERENCES

- Albert, A. and J. A. Andersen (1984). On the existence of maximum likelihood estimates in logistic regression models. *Biometrika* 71(1), 1–10.
- Bang, H. and J. M. Robins (2005). Doubly robust estimation in missing data and causal inference models. *Biometrics* 61(4), 962–973.
- Cao, W., A. A. Tsiatis, and M. Davidian (2009). Improving efficiency and robustness of the doubly robust estimator for a population mean with incomplete data. *Biometrika*, 723–734.
- Cassel, C. M., C. E. Sarndal, and J. H. Wretman (1976). Some results on generalized difference estimation and generalized regression estimation for finite populations. *Biometrika* 63(3), 615–620.
- Cover, T. M. and J. A. Thomas (2012). *Elements of information theory*. John Wiley & Sons.
- Diamond, A. and J. S. Sekhon (2013). Genetic matching for estimating causal effects: A general multivariate matching method for achieving balance in observational studies. *Review of Economics and Statistics* 95(3), 932–945.
- Drake, C. (1993). Effects of misspecification of the propensity score on estimators of treatment effect. *Biometrics*, 1231–1236.
- Ferwerda, J. (2014). Electoral consequences of declining participation: A natural experiment in austria. *Electoral Studies* 35(0), 242–252.
- Freedman, D. A. and R. A. Berk (2008). Weighting regressions by propensity scores. *Evaluation Review* 32(4), 392–409+.
- Hahn, J. (1998). On the role of the propensity score in efficient semiparametric estimation of average treatment effects. *Econometrica* 66(2), 315–332.
- Hainmueller, J. (2011). Entropy balancing for causal effects: A multivariate reweighting method to produce balanced samples in observational studies. *Political Analysis*, mpr025.

- Hirano, K. and G. Imbens (2001). Estimation of causal effects using propensity score weighting: An application to data on right heart catheterization. *Health Services and Outcomes Research Methodology* 2(3-4), 259–278.
- Hirano, K., G. W. Imbens, and G. Ridder (2003). Efficient estimation of average treatment effects using the estimated propensity score. *Econometrica* 71(4), 1161–1189.
- Holland, P. W. (1986). Statistics and causal inference. *Journal of the American Statistical Association* 81, 945–960.
- Imai, K., G. King, and E. A. Stuart (2008). Misunderstandings between experimentalists and observationalists about causal inference. *Journal of the Royal Statistical Society: Series A (Statistics in Society)* 171(2), 481–502.
- Imai, K. and M. Ratkovic (2014). Covariate balancing propensity score. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)* 76(1), 243–263.
- Kang, J. D. and J. L. Schafer (2007). Demystifying double robustness: A comparison of alternative strategies for estimating a population mean from incomplete data. *Statistical Science* 22(4), 523–539.
- Lunceford, J. K. and M. Davidian (2004). Stratification and weighting via the propensity score in estimation of causal treatment effects: a comparative study. *Statistics in Medicine* 23(19), 2937–2960.
- Marcus, J. (2013). The effect of unemployment on the mental health of spouses evidence from plant closures in germany. *Journal of Health Economics* 32(3), 546–558.
- Qin, J. and B. Zhang (2007). Empirical-likelihood-based inference in missing response problems and its application in observational studies. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)* 69(1), 101–122.
- Ridgeway, G. and D. F. McCaffrey (2007). Comment: Demystifying double robustness: A comparison of alternative strategies for estimating a population mean from incomplete data. *Statistical Science* 22(4), 540–543.
- Robins, J., M. Sued, Q. Lei-Gomez, and A. Rotnitzky (2007). Comment: Performance of double-robust estimators when inverse probability weights are highly variable. *Statistical Science* 22(4), 544–559.
- Robins, J. M., A. Rotnitzky, and L. Zhao (1994). Estimation of regression coefficients when some regressors are not always observed. *Journal of the American Statistical Association* 89, 846–866.
- Robins, J. M. and N. Wang (2000). Inference for imputation estimators. *Biometrika* 87(1), 113–124.
- Rosenbaum, P. and D. Rubin (1983). The central role of the propensity score in observational studies for causal effects. *Biometrika* 70, 41–55.
- Rosenbaum, P. and D. Rubin (1984). Reducing bias in observational studies using subclassification on the propensity score. *Journal of the American Statistical Association* 79, 516–524.
- Rosenbaum, P. R. et al. (2002). Covariance adjustment in randomized experiments and observational studies. *Statistical Science* 17(3), 286–327.
- Rothe, C. and S. Firpo (2013). Semiparametric estimation and inference using doubly robust moment conditions. IZA Discussion Papers 7564, Institute for the Study of Labor (IZA).

- Rotnitzky, A., Q. Lei, M. Sued, and J. M. Robins (2012). Improved double-robust estimation in missing data and causal inference models. *Biometrika* 99(2), 439–456.
- Rubin, D. (1974). Estimating causal effects of treatments in randomized and non-randomized studies. *Journal of Educational Psychology* 66(5), 688–701.
- Rubin, D. B. (1976a). Multivariate matching methods that are equal percent bias reducing, i: Some examples. *Biometrics* 32(1), pp. 109–120.
- Rubin, D. B. (1976b). Multivariate matching methods that are equal percent bias reducing, ii: Maximums on bias reduction for fixed sample sizes. *Biometrics* 32(1), pp. 121–132.
- Sekhon, J. S. (2011). Multivariate and propensity score matching software with automated balance optimization: The Matching package for R. *Journal of Statistical Software* 42(7), 1–52.
- Silvapulle, M. J. (1981). On the existence of maximum likelihood estimators for the binomial response models. *Journal of the Royal Statistical Society. Series B (Methodological)* 43(3), 310–313.
- Smith, J. A. and P. E. Todd (2001). Reconciling conflicting evidence on the performance of propensity-score matching methods. *American Economic Review* 91(2), 112–118.
- Stefanski, L. A. and D. D. Boos (2002). The Calculus of M-Estimation. *The American Statistician* 56(1), 29–38.
- Tan, Z. (2006). A distributional approach for causal inference using propensity scores. *Journal of the American Statistical Association* 101, 1619–1637.
- Tan, Z. (2010). Bounded, efficient and doubly robust estimation with inverse weighting. *Biometrika* 97(3), 661–682.
- Wang, Q. and J. N. K. Rao (2002). Empirical likelihood-based inference under imputation for missing response data. *The Annals of Statistics* 30(3), 896–924.